



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Pneumonia.—Pneumonia deaths numbered 241, a rate of 116.8 per 100,000. In the corresponding month last year the rate was 87.7. The disease prevailed very much more extensively in the cities than in the country, the rates being, respectively, city 179.1 per 100,000, country 84.5. Pneumonia does not lead phthisis pulmonalis as a cause of death in Indiana, and has only done so in one or two months in the last four years.

MASSACHUSETTS—*Worcester*.—Month of October, 1903. Estimated population, 123,331. Total number of deaths, 148, including diphtheria 1, enteric fever 2, scarlet fever 1, and 20 from tuberculosis.

Month of November, 1903. Total number of deaths, 129, including enteric fever 1, and 14 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended December 26, 1903, from 78 observers indicate that scarlet fever, erysipelas, intermittent fever, smallpox, whooping cough, and dysentery were more prevalent, and diarrhea, diphtheria, measles, and meningitis were less prevalent than in the preceding week.

Meningitis was reported present at 3, whooping cough at 8, measles at 29, diphtheria at 37, enteric fever at 56, smallpox at 58, scarlet fever at 80, and phthisis pulmonalis at 184 places.

OHIO—*Cincinnati*.—Month of November, 1903. Estimated population, 336,000. Total number of deaths, 481, including diphtheria 6, enteric fever 9, measles 7, scarlet fever 5, smallpox 2, and 57 from tuberculosis.

WISCONSIN—*Milwaukee*.—Month of November, 1903. Estimated population, 315,000. Total number of deaths, 325, including diphtheria 5, enteric fever 4, scarlet fever 1, and 34 from tuberculosis.

Report of immigration at Baltimore.

OFFICE OF THE COMMISSIONER OF IMMIGRATION,
Port of Baltimore, Md., January 2, 1904.

Number of alien immigrants who arrived at this port during the week ended January 2, 1904; also names of vessels and ports from which they came.

Date.	Vessel.	Where from.	Number of immigrants.
Dec. 29	Brandenburg.....	Bremen.....	751
29	Hippolyte Dumois.....	Ceiba, Honduras.....	1
31	Koln.....	Bremen.....	417
	Total.....		1,169

LOUIS T. WEIS, *Commissioner*.